

a living river

CHARTING SANTA CRUZ RIVER CONDITIONS
NORTHWEST TUCSON TO MARANA—2018 WATER YEAR



SUPPLEMENTARY REPORT FOR 2013 TO 2018 WATER YEARS



SONORAN
INSTITUTE



PIMA COUNTY

THE SANTA CRUZ RIVER

A LIVING ECOSYSTEM



The Santa Cruz River in northwest Tucson and Marana flows year-round and provides the principal wetland habitat in Pima County. River flows are sustained by the release of effluent—highly-treated wastewater—from two regional facilities. In December 2013, Pima County completed the largest public works project in southern Arizona by investing over \$600 million to upgrade the treatment process. Improved treatment affords the opportunity to enhance the aquatic environment along the river, reduce odors, and increase re-use of reclaimed water.

The *Living River* reports were developed to annually gauge conditions of this valuable ecosystem and track the impacts of our community investment. This supplementary report summarizes data from six years (2013–2018 water years). The pages following this executive summary provide details on the water context and data for 16 indicators of river health. A technical committee of experts selected these indicators as described in a selection process report.

All *Living River* reports can be found on the Sonoran Institute website at www.sonoraninstitute.org

CHANGES IN WATER QUALITY AND WETLAND CONDITIONS

- **Ammonia no longer limiting life:** Ammonia, which can be toxic to aquatic organisms, was appreciably reduced to low levels.
- **Oxygen availability not a stressor:** Essential for aquatic life, dissolved oxygen remained at steady levels or increased. Biochemical oxygen demand (an indirect measure of pollutants that use up oxygen in the water) declined to nearly non-detectable levels, indicating that there is more oxygen available for organisms to thrive.
- **Water clarity much improved:** Sediments and other particles carried in the water decreased, resulting in clear river water on normal non-flooding days. Elevated sediment levels in the water can increase water temperature, thereby decreasing available dissolved oxygen.
- **More diverse life:** Improvements in water quality has allowed aquatic life in the river to rebound. Six species of fish, including the endangered Gila topminnow, and increased diversity of aquatic invertebrates (which include insects, crustaceans, and worms) have been observed.
- **Reduced flow extent:** The length of the flowing river has decreased and is more variable due to a combination of factors, including increased water infiltration from reduced nutrient levels, scouring floods, reductions in volume of water released, and changes in flow management.
- **Wetland plants reduced in drying sections:** The release of effluent supports wetland plants and trees. There is a decrease in willows and increased variability in streamside plants in the sections of reduced flow extent.
- **Very little odor escaped the reclamation facility boundary:** Odor levels are far below levels required by facility permits and anecdotal observations report odor as hardly noticeable near the facility boundaries.



Cloudy water, before upgrade



Clear water, after upgrade



Gila topminnow, *Poeciliopsis occidentalis*

OTHER OBSERVATIONS







- **Total effluent released to the river has decreased:** Releases of effluent have decreased an average of 13% since 2013, with lowest volume released in 2018. However, effluent remains the primary source of water in the river. Stormwater is also an important source of flows and total volume of stormwater in the river has increased since 2013.
- **Increased infiltration rates and groundwater recharge:** The amount of water that recharged local aquifers more than doubled between 2013 and 2018. This is likely from increased rates of infiltration resulting in part from improved water quality.
- **Many kids are seeing a flowing river for the first time:** The Living River of Words youth art and science program continues to provide the first contact with a flowing stream for hundreds of kids. The Santa Cruz River from northwest Tucson to Marana provided meaningful inspiration for youth art and poetry projects. To date over 3,000 youth have visited the river since 2015.

ASSESSING CONDITIONS

The Living River report evaluates conditions of the Santa Cruz River from northwest Tucson to Marana using indicators (see table below) organized into six categories that represent a breadth of biological, chemical, physical, and social properties

of the river. The indicators relate to conditions in the river channel and in the riparian areas, the areas next to and affected by the river.

The purpose of the Living River series is to monitor and report on wetland and riparian conditions at various intervals downstream of the effluent discharge points. As effluent flows downstream, it impacts and is impacted by the natural

| CATEGORY | | PURPOSE | INDICATORS |
|-----------------------------|---|--|--|
| Flow Extent |  | Water flowing in and out of the system determines available aquatic habitat. | <ul style="list-style-type: none"> • Miles of flow in June • Number of “dry days” at Trico Road |
| Water Clarity |  | Solid particles in the water and on the riverbed can impact habitat and conditions for aquatic life. | <ul style="list-style-type: none"> • Total suspended solids • Turbidity • Percent fines on riverbed |
| Water Quality |  | Specific chemical conditions are necessary to sustain the river's animal and plant communities. | <ul style="list-style-type: none"> • Total dissolved solids • Ammonia • Dissolved oxygen • Biochemical oxygen demand • Metals |
| Aquatic Wildlife |  | Wildlife in the river integrate and reflect conditions of many factors of the surrounding environment. | <ul style="list-style-type: none"> • Fish • Aquatic invertebrates |
| Riparian vegetation* |  | Plant communities reflect changes in water quantity and quality. | <ul style="list-style-type: none"> • Wetland indicator status • Nitrogen affinity score • Riparian tree cover |
| Social Impacts |  | Aesthetic factors directly impact people living or recreating along the river. | <ul style="list-style-type: none"> • Odor at reclamation facilities |

conditions of soils, vegetation, and the surrounding ecosystem. For the purposes of this study, the 23-mile stretch of river is divided into three sections, or reaches: Three Rivers, Cortaro Narrows, and Marana Flats. Reaches were delineated by their differing hydrology, geology, and adjacent land use.

Data are collected and summarized by water year (October 1–September 30) and compared to the baseline conditions observed in the 2013 water year. This supplemental report shares data from all the water years to enable an easy viewing of trends and long-term patterns.

*Riparian vegetation only monitored from 2013–2016


Streamflow, Rainfall, and Water Budget

Streamflow, or the amount of water flowing in a river, provides an important context for the results of the indicators.

Reclamation facilities continuously release water into the river, which accounts for the majority of daily streamflow.

However, streamflow also includes stormwater, which is influenced by rainfall and the amount of impervious area (e.g.,

roadways) in the watershed. The Santa Cruz River Watershed includes all of the land whose stormwater flows toward the river. Seasonal floods are important for recharging aquifers, dispersing seeds, inducing seed germination, and clearing natural debris.



2013–2018 RAINFALL

Rainfall totals from the Tucson International Airport (TIA) and near the river at Cortaro Farms Road (CFR) provide a general idea of when stormwater may have increased streamflow.

TIA had an annual average of 11 inches of rain. The most rain fell in 2016 and 2015 with 13 and 14 inches respectively. The historical average is 11 inches from 1949 to 2011.

- The winter rains ranged from 1 to 6 inches of rain.
- The summer monsoon ranged from 2 to 9 inches of rain.

CFR had an average of 11 inches of rain. The most rain fell in 2016 and 2018 with 12 and 14 inches respectively. This station was set up in 2012 and has no historical data.

- The winter rains ranged from 1 to 5 inches of rain.
- The summer monsoons ranged from 3 to 10 inches of rain.

2013–2018 STREAMFLOW


Streamflow is measured with gages at Cortaro Road and Trico Road, which are downstream of the Agua Nueva and Tres Rios Reclamation Facilities. Streamflow, measured in cubic feet per second (cfs), is the volume (cubic feet) of water flowing past a fixed point in a specific time period (1 second). Measuring streamflow daily tracks seasonal floods.

At Cortaro Road, mean daily streamflow has remained similar over the years, though flood peaks have been higher since 2013. Flows at Trico Road has become more variable and have declined since the facility upgrades were complete in December of 2013. Like Cortaro Road, the peak flows have increased. Between 2015 and 2017 water years, the river is dry at Trico Road between rainy seasons; this was not the case in 2018 where there are fewer dry periods.

Streamflow, Rainfall, and Water Budget, cont.

A water budget estimates the water inputs and outputs. Inputs are effluent and stormwater, while outputs include water that does one of the following: flows past Trico Road (the end of the study area), evaporates or is used by wetland vegetation (a process called evapotranspiration), is diverted for agricultural use, or sinks into the riverbed to recharge local groundwater. Volumes are totaled in acre-feet (AF), the

number of acres that would be covered with water one foot deep. Total recharge volume is for effluent only and does not include stormwater. Totals for evapotranspiration and diversions are not directly measured. They are estimates and viewed as fairly constant from year to year in the calculations for the managed recharge projects located along the Lower Santa Cruz River.



* Includes effluent that doesn't go to the river; either it is diverted to the reclaimed system for irrigation or to recharge basins located outside the river channel.

¹ Excluding days with stormwater, the volume of only effluent flowing past Trico Road is: 2013 = 26,800 AF; 2014 = 13,400 AF; 2015 = 2,100 AF; 2016 = 3,800 AF; 2017 = 3,700 AF; 2018 = 6,000 AF

2013–2018 WATER BUDGET


Effluent provides most of the flow in the river. Effluent releases from Agua Nueva have decreased for two reasons. First, the facility upgrade completed in December of 2013 resulted in some wastewater being redirected to Tres Rios and released further downstream. Second, more effluent was diverted, before release into the river, into nearby basins to recharge local aquifers. Diverted effluent reached a steady volume starting in 2017 water year. Overall, effluent inputs were reduced by an average of 13% since 2013, with 2018 representing the lowest volume released. Total inputs increased with higher volumes of stormwater. Even with greater inputs, recharge increased significantly, likely from higher infiltration rates after improved water quality. Increased infiltration has also reduced the amount of water that flows past Trico Road. Exact volumes of water diverted for agriculture and used by wetland vegetation are not known and treated as constants in recharge calculations.



FLOW EXTENT

Measuring flow extent, or the distance the river is flowing, is a quick visual way to track changes in water inputs and outputs, while providing a rough measure of the quantity of aquatic habitat available. For example, high flow extent may indicate high inputs and high availability of habitat for aquatic life. Low flow extent may indicate reduced inputs, which could decrease aquatic habitat. Alternatively, low flow extent could indicate greater recharge of water into local aquifers.

Miles of flow to Trico Road in June in each reach prior to the monsoon season determines the minimum extent of flow during the driest time of year. This is typically measured on one morning in mid-June. **Flow at Trico Road**, located at the farthest end of the study area, estimates daily changes in maximum flow extent through the three reaches by counting the “dry days,” or days with no streamflow.



2013–2018 RESULTS

Flow extent decreased and was more variable after the December 2013 upgrades. In June 2013, the river flowed uninterrupted to the end of the 23 mile study area, and continued another 5 miles further into Pinal County. Since 2013, only Cortaro Narrows flowed through the whole reach in June. Though variable in length, dry stretches of the river formed between Agua Nueva and Tres Rios, and close to Trico Road.


Reduced flow extent is primarily due to increased recharge. However, water management is also an important factor and has contributed to the drying of the Three Rivers reach. Releases from Agua Nueva have decreased for two reasons. First, the upgrade resulted in some wastewater being redirected to Tres Rios and thus released further downstream. Second more water has been diverted to recharge basins adjacent to the river. In Marana Flats, water is also diverted by an earthen berm for irrigation of agriculture and for recharge at Marana High Plains, a constructed recharge basin adjacent to the river. The berm failed just before the June 2016 survey, and may have increased miles of flow recorded that year.

The daily flow at Trico Road has become more variable with increased dry days where there is no water in the river. In addition to increased recharge, natural and human management of river flow has likely influenced conditions at Trico Road.



FLOW EXTENT: Continued

In September 2014, a large flood moved the location of the low-flow channel and breached a berm along the El Rio Preserve, a former borrow pit near the start of Marana Flats. This allowed water to flow into the pit and form the wetlands at El Rio Preserve. The river stopped flowing into the wetlands in January 2015 when a flood moved the low-flow channel again, demonstrating nature's contribution to water management. The scouring flood may have further increased infiltration rate. This combined with diversion of flow into the wetlands may have increased the number of dry days recorded in 2015. Conversely, a berm diverting water for agriculture and recharge at Marana High Plains failed several times in 2016 may have decreased the dry days at Trico Road. Recently, number of dry days has decreased, even as total water in the river has decreased.






WATER CLARITY: Total Suspended Solids

Rivers naturally move sediments and other small particles of algae or detritus downstream. High concentrations of materials in the water can create murky “dust storm” conditions and may impact conditions for aquatic life. Under chronically high “dust storm” conditions, sunlight doesn’t travel as deep into the water. Thus, aquatic plants may not receive enough sunlight to conduct photosynthesis and

aquatic predators may not be able to see well enough to capture prey.

Total suspended solids is an estimate of the number of particles in the water, or the intensity of the “dust storm.” ADEQ does not have a standard for total suspended solids. The results from the 2013 water year serve as a baseline.



2013–2018 RESULTS


Total suspended solids (TSS) was measured a total of 87 times. Levels of TSS decreased in Cortaro Narrows and Marana Flats after the upgrades were complete. Levels of TSS were similar in all three reaches in 2014 and 2015. Since 2016, Marana Flats has increased, though levels were still lower than the 2013 baseline. Stormwater is expected to have greater levels of TSS. Samples of stormwater are collected upstream of Agua Nueva when possible. Four samples collected (one each year during the summer monsoon for 2013–2016) had TSS concentrations ranging from 1,050 to 46,300 mg/L and were higher than levels on normal flow conditions.



WATER CLARITY: Turbidity

Rivers naturally move sediments and other small particles of algae or detritus downstream. High concentrations of materials in the water can create murky “dust storm” conditions and may impact conditions for aquatic life. Under chronically high “dust storm” conditions, sunlight doesn’t travel as deep into the water. Thus, aquatic plants may not receive enough sunlight to conduct photosynthesis, and aquatic predators may not be able to see well enough to

capture prey. **Turbidity** measures water clarity, or how far you can see through the “dust storm,” and is reported in Nephelometric Turbidity Units (NTU). High NTU indicates the water is cloudy and hard to see through. The 1988–1993 median level of turbidity in the Cortaro Narrows reach was 15 NTU. ADEQ does not have a standard for turbidity, so this assessment uses 15 NTU as a historical reference value.



2013–2018 RESULTS


Turbidity was measured throughout the year at several locations for a total of 92 times. Overall, the reference value was met 72 times (78%). Average turbidity within each reach has decreased since the 2013 baseline. Although turbidity decreased in Marana Flats following the upgrades, values have been higher and most variable in this reach since 2016.



WATER CLARITY: Percent Fines

Rivers naturally move sediments and other small particles of algae or detritus downstream. High concentrations of materials in the water can create murky “dust storm” conditions and may impact conditions for aquatic life. Under chronically high “dust storm” conditions, sunlight doesn’t travel as deep into the water. Thus, aquatic plants may not receive enough sunlight to conduct photosynthesis, and aquatic predators may not be able to see well enough to capture prey.

Percent fines is an estimate of the portion of the riverbed comprised of small sediments (≤ 2 mm in diameter). Fines, or “muck,” that settle out of the storm onto the riverbed can become so abundant that they smother aquatic life and habitat. ADEQ does not have a standard for rivers dominated by effluent. This assessment uses the reference value for warm-water rivers, $<50\%$.



2013–2018 RESULTS


Percent fines were estimated at four sites where macroinvertebrate samples were collected. Overall there was a reduction in the percent fines covering the riverbed at these sites, though there was a lot of variation. Due to reductions in flow extent, the second survey site in Three Rivers and the survey site in Marana Flats had to be shifted upstream in 2015 and 2014 respectively.



WATER QUALITY: Total Dissolved Solids

Many of the dissolved solids are essential nutrients for plants and animals, but when too abundant they can produce unhealthy conditions for aquatic life and riparian vegetation. Thus, measuring **total dissolved solids** (TDS) is commonly used to monitor excess salts in the water. TDS in the effluent has been rising since the 1990s with increased use of

Colorado River water in the Tucson area. The Colorado River has greater TDS, mostly in form of dissolved salts, than the local groundwater. Because there is no standard for TDS (often standards are for individual elements that contribute to TDS), the results from the 2013 water year will serve as a baseline.



2013–2018 RESULTS


Total dissolved solids (TDS) were measured 91 times. Overall, levels of TDS were similar in all three reaches. Generally TDS hasn't changed very much, though variability in TDS levels decreased in 2015 and 2016 for unknown reasons. The lowest measure of TDS was in Marana Flats. This sample was collected on a day where there was possible stormwater influence. Thus, the addition of water with lower TDS levels may have diluted the levels in this reach of the Santa Cruz River. Samples of stormwater are collected upstream of Agua Nueva when possible. Four samples collected (one each year during the summer monsoon for 2013–2016) averaged 280 mg/L. No stormwater sample was collected in 2017 or 2018.



WATER QUALITY: Ammonia

Nitrogen is an essential nutrient for plant and animal life, but too much can contribute to nutrient pollution. Nutrient pollution, such as high levels of nitrogen and phosphorus, enters the river from air pollution, fertilizer, surface runoff, and the release of effluent. While elevated nutrient levels can benefit riparian plants, they can also lead to poor water quality conditions for aquatic wildlife.

Ammonia (NH_3) is one form of nitrogen that can be toxic to fish. Even at low concentrations, ammonia can reduce hatching success, among other impacts. The ADEQ standard for ammonia varies with pH (level of acidity) and temperature. As pH and temperature increase, the toxicity of ammonia increases; thus, the acceptable level of ammonia decreases with high pH and temperature.



Upgrades to reclamation facilities complete (Dec 2013)

Note - Ammonia standards vary with temperature and pH and can't be graphed as a single line

2013–2018 RESULTS


Ammonia was measured 90 times along the river. Overall the standard was met 63 of the 90 times (70%). Levels of ammonia have dropped significantly after the upgrade was complete in 2013. Levels of ammonia also decreased with distance from the reclamation facilities, as it breaks down into other forms of nitrogen while moving downstream. Although there were exceedances of the ammonia standard before and after the upgrade, the magnitude of the average exceedance has dropped significantly (average exceedance above standard: *before upgrade* = 14 mg/L, *after upgrade* = 1.6 mg/L). The percent of samples with elevated levels of ammonia has also decreased (*before upgrade* = 75% of samples, *after upgrade* = 20% of samples). Nitrogen removal is a complex process. Pima County Regional Wastewater Reclamation Department is evaluating system processes at Agua Nueva and installing new technologies at Tres Rios in 2019 to address occasional, short-lived periods of elevated ammonia in the river.



WATER QUALITY: Dissolved Oxygen

Fish and other aquatic animals need **dissolved oxygen** to survive. Rivers absorb oxygen from the atmosphere, and aquatic plants and algae produce oxygen. Natural causes of variability in dissolved oxygen levels include nutrient

levels, shading, water temperature, and time of day. ADEQ sets the minimum standard for dissolved oxygen in streams dominated by effluent at 3 milligram per liter (mg/L) during the day (3 hrs after sunrise to sunset).



2013–2018 RESULTS


Dissolved oxygen was measured 92 times along the river. All of the samples met the standard for dissolved oxygen (100%). Levels of dissolved oxygen stayed fairly constant in Three Rivers and Cortaro Narrows. However, Marana Flats saw an increase in dissolved oxygen after the facility upgrades were completed.



WATER QUALITY: Biochemical Oxygen Demand

Biochemical oxygen demand (BOD) is an estimate of how much dissolved oxygen is being used. Microorganisms in the river consume dissolved oxygen as they break down organic materials such as leaves and woody debris, dead plants and animals, and animal wastes. If there are a lot of organic materials in the water, these microorganisms become

so numerous that they consume much of the dissolved oxygen and deprive other aquatic animals of the oxygen they need to survive. Though there are standards for BOD in the wastewater reclamation process, there is no standard for BOD in rivers. The results from the 2013 water year will serve as a baseline.



2013–2018 RESULTS

Biochemical oxygen demand was measured 92 times along the river. BOD has decreased since the upgrades to the reclamation facilities were completed. The high levels observed in Cortaro Narrows are absent after the 2013 water year. This pattern is generally the same in Marana Flats. However, for reasons unknown, measures of BOD in the first half of 2017 were similar to the high levels observed during the 2013 baseline.



WATER QUALITY: Metals

Metals in high concentrations endanger wildlife in aquatic ecosystems by lowering reproductive success, interfering with growth and development, and, in extreme cases, causing death. Most metals build up in aquatic food chains and may pose long-term threats to all organisms in the aquatic environment. Rivers are exposed to pollutant

metals through numerous sources, including mine drainage, roadways, and by the release of metals naturally occurring in near-surface rocks and sediments. ADEQ has set standards for the protection of aquatic wildlife. Results for the following metals are compared to their appropriate standard: arsenic, cadmium, chromium, copper, lead, mercury, and zinc.

Average values for dissolved metals tested throughout the year
concentrations in micrograms/liter (ug/L), also known as parts per billion (ppb)

Average Standard
standards for wildlife vary
with water hardness

| | Direction of Flow | | | | |
|----------|-------------------|-----------------|--------------|-----|------------|
| | Agua Nueva >> | | Tres Rios >> | | |
| Arsenic | 3.9 | 3.0 | 3.2 | 3.5 | 150 ug/L* |
| Cadmium | ND | ND | ND | ND | 2.9 ug/L |
| Chromium | 0.7 | 0.6 | 0.5 | 0.5 | 11 ug/L* |
| Copper | 2.4 | 2.2 | 2.3 | 2.3 | 18 ug/L |
| Lead | 0.3 | 0.2 | 0.3 | 0.4 | 6 ug/L |
| Mercury | ND | ND | ND | ND | 0.01 ug/L* |
| Zinc | 50 | 48 | 45 | 38 | 236 ug/L |
| | Three Rivers | Cortaro Narrows | Marana Flats | | |

*set value, not an average

ND = Not Detected

2013–2018 RESULTS


All samples tested over the six years have met the appropriate standard for the following dissolved metals: arsenic, cadmium, chromium, copper, lead, mercury, and zinc. The samples taken within Marana Flats were from three different sites and averaged here. The sample location had to be moved several times due to drying and inconsistent flows following increased recharge rates.



AQUATIC WILDLIFE: Fish

Fish can serve as effective indicators of river health because they live for several years and vary in their tolerance to pollution. Historically, the Santa Cruz River supported several native fish species: Gila topminnow, Gila chub, desert sucker, Sonora sucker, longfin dace, and a pupfish species that went

extinct when the river ceased to flow year-round. There is no standard for abundance or diversity of fish. The results from the 2013 water year will serve as a baseline for measuring change in subsequent years.



2013–2018 RESULTS

Fish surveys were conducted annually in the fall at the four locations macroinvertebrates were surveyed. Surveys aim to detect all fish species present at a location, but do not try to assess population numbers. Improvements in water quality have allowed fish to thrive. Overall, number of fish species observed increased from one to six. All are non-native, except for the endangered Gila topminnow, which was found in the river in 2017. Exactly how this native fish returned is unknown. Gila topminnow are found in the Santa Cruz near Nogales and may have been carried in flood flows down to the Three Rivers reach. Although flood flows connected these stretches of Santa Cruz River for two days in July 2017, initial genetic analysis suggests the Gila topminnow in the river near Tucson are most similar to fish found in the Cienega Creek watershed.


Recording the most species, Cortaro Narrows may provide the most diverse habitat for fish. Flows in Three Rivers are often very shallow and may favor smaller fish like the Western mosquitofish and Gila topminnow. Though presence of large fish, likely common carp, was reported in the media in spring of 2014, larger species were not observed or captured until the 2015 survey. A large 20,000 cfs flood in September 2014 may have washed them past Trico Road and beyond the study area. Flood conditions (around 4,000 cfs) two weeks prior to the 2018 survey may have also influenced number of species observed.



AQUATIC WILDLIFE: Aquatic Invertebrates

Aquatic invertebrates break down organic materials and are important prey for fish and other species. They also differ in their tolerances to pollution. Chironomidae (midges) are pollution tolerant and found in high numbers even with low oxygen levels and high organic matter. Ephemeroptera (mayflies) have exposed gills on the outside of their body, making them very pollution sensitive. The percent of the invertebrate community comprised of Ephemeroptera taxa

is commonly used to help track changes in water quality. Regardless of sensitivity to pollution, if a single species or group accounts for more than 50% of the community, this lack of diversity suggests a stream is impaired. Another way to look at diversity is simply looking at the total number of different invertebrate taxa found in the samples collected.



2013–2018 RESULTS


The aquatic invertebrate community was surveyed annually at the four locations that fish were surveyed. Invertebrates were sampled using the standard operating procedure developed by Arizona Department of Environmental Quality which involves kick-net samples in riffles. This gives a quick assessment of the site's biological integrity. Overall, there were signs of improvement. Diversity was higher after the upgrades were complete, with the number of taxa at each site increasing. Increased diversity is also demonstrated by the smaller percentage of the community composed of a single group. If the dominant group is more than 50% of the community, river life is thought to be impaired. Improvements are also supported by the increase in the percent of the community comprised of pollution-sensitive species from the order Ephemeroptera. The 2017 and 2018 water years reached the highest levels of diversity across most sites with the highest number of taxa recorded and the lowest recorded percentage of the dominant group.



RIPARIAN VEGETATION: Wetland Indicator Status

Wetland indicator status measures abundance of stream-side plants that vary in their need for permanent water in the river channel. Scores range from 1 to 5. Low scores (<4) indicate that the majority of plants at a given location are wetland plants like watercress and cattails, which depend on consistent presence of water in the river. High scores

(>4) indicate that the majority of plants are upland plants like burrobrush and different grasses; these do not depend on consistent presence of water in the river and usually are not found in wetlands. Results from the 2013 water year will serve as a baseline to help track future changes in wetland plants.



2013–2016 RESULTS


Not measured in 2017 or 2018. Wetland indicator status (WIS) was determined for eight total locations along the river. Overall, scores have remained similar at most sites. Scores averaged 2.7 downstream of Agua Nueva. This suggests greater presence of wetland plants instead of upland plants as the river flowed away from the reclamation facilities. Just upstream of the study area, a reference site had the highest scores and was dominated by upland plants. Two sites (approximately 5 and 20 miles downstream) appeared to shift toward more upland plants with increased scores in 2015. This may be in part explained by changes in flow extent, as these sites experienced dry conditions more frequently in water year 2015. However, these same sites were wet again when surveyed in 2016, and stream-side plants shifted back toward wetland plants.



RIPARIAN VEGETATION: Nitrogen Affinity Score

Although nitrogen is an essential nutrient, too much can undermine plant growth or favor the growth of plants that thrive in high-nitrogen environments. **Nitrogen affinity score** measures the abundance of stream-side plants that vary in their tolerance of nitrogen. Scores range from 1 to 9. Low scores (<5) indicate that the majority of plants at a given location grow well with low levels of nitrogen, like burrobrush

and different grasses. High scores (>5) indicate that the majority of plants grow well with high levels of nitrogen, like cattails and common sunflowers. Changes in nitrogen affinity scores likely reflect changes in water quality, either an increase or decrease in nutrients in the water. Results from the 2013 water year will serve as a baseline.



2013–2016 RESULTS


Not measured in 2017 or 2018. Nitrogen affinity score was determined for eight total locations along the river. Overall, scores have remained similar at most sites. Scores averaged 6.2 downstream of Agua Nueva. This suggests that stream-side plants that grow well in high nitrogen environments were most common immediately downstream of the reclamation facilities. Just upstream of the study area a reference site had the lowest scores and was dominated by plants that grow well with low levels of nitrogen. Two sites (approximately 5 and 20 miles downstream) appeared to shift toward more low-nitrogen plants in 2015. Though we may expect this shift from reduced nutrient pollution, reduction in water presence and soil moisture may be the bigger factor. Both of these sites experienced dry conditions more frequently in water year 2015 and were dry at time of survey in 2015. These same sites shifted back towards nitrogen-loving plants in 2016, when water was present again at time of survey. So both nitrogen affinity and wetland indicator seem to indicate presence of permanent water in the channel or high soil moisture. This is supported by a high correlation of the nitrogen scores with wetland scores; plants with high nitrogen scores had very low wetland scores, or more simply, the wetland plants in our area love nitrogen.



RIPARIAN VEGETATION: Riparian Tree Cover

Riparian tree cover measures the abundance of adult trees along the river and in the adjacent floodplain. High tree cover indicates the presence of sufficient soil moisture to support riparian trees. Tree cover is commonly reported as basal area. Basal area, measured in square meters per hectare (m^2/ha), is the area covered by trees in one hectare (10,000 m^2 , or approximately two football fields). In addition, riparian tree species differ in their tolerance to declines in soil moisture. Native cottonwoods and willows have shallow roots

and are more sensitive to reductions in soil moisture. Velvet mesquite and non-native tamarix species, such as Athel tamarix and saltcedar, have deeper roots and can tolerate a greater range of soil moisture. Trees grow slowly, and amount of cover is not likely to change on an annual basis, unless vegetation is affected by sustained drying or large floods. Tree cover was measured in 2015, and results from the 2013 water year serve as a baseline.



2013–2015 RESULTS


Only measured in 2013 and 2015. Overall tree cover, as measured by basal area and percent canopy cover, decreased between 2013 and 2015 (trees were not measured in 2016). Most notable was the decrease in cover of Goodding's willow. Decrease in cover of mature trees is likely the result of decreased flow extent. There may not have been enough moisture to support more shallow rooted trees like Goodding's willow. More monitoring will be needed to determine if effluent continues to support mature riparian trees in all three reaches, and whether the community shifts to deeper rooted trees such as velvet mesquite and non-native tamarix species.



SOCIAL IMPACTS: Odor at the Reclamation Facilities

Reclamation facilities are restoring a piece of the river heritage and supporting important wetland habitats by releasing effluent into the river. However, unpleasant odors often associated with the reclamation process can lead to negative perceptions of the river for those living near or recreating along the river. The most common offender is

hydrogen sulfide (H_2S) or the “rotten egg” smell. **Odor at the reclamation facilities** tracks changes in odors linked to the reclamation process. Minimizing both the extent and intensity of disagreeable odors coming from the facilities was one of the goals of the reclamation facility upgrades.



2013–2018 RESULTS

Initial facility improvements in 2007 reduced both the extent and intensity of odor emanating from the reclamation facilities. Data to map extent and intensity of odors since 2013 is not available. However, as of 2016, odor is monitored continuously at the facilities and at numerous points along the surrounding fencelines. Between 2016 and 2018 there were no odor complaints at Agua Nueva and levels of hydrogen sulfide (H_2S) never exceeded 10 parts per billion (ppb), the level allowed by facility permits. Levels of H_2S at Tres Rios were also low with an average of 0.1 ppb for over 11 million measures of odor between 2016 and 2018. Levels of odor exceeded the 10 ppb permit level less than 0.3% of the time. Detailed odor data of this kind is not available for years prior to the upgrades, thus comparisons to previous H_2S levels are not possible.

ACKNOWLEDGEMENTS

Sonoran Institute and Pima County prepared this report with generous funding from Pima County Regional Wastewater Reclamation Department, Pima County Regional Flood Control District, and community individuals. We are grateful for the expert guidance from our Living River Technical Committee, and for the support of our project partners, including Arizona Department of Environmental Quality, Arizona State University, Tucson Audubon Society, University of Arizona, and the U.S. Geological Survey.

The Sonoran Institute convened a Living River Technical Committee of ecology, hydrology, and wildlife experts to bring the best available science to bear on the development of the *Living River* health assessments. The Technical Committee provided guidance by selecting and aggregating indicators of river health, identifying reference values or standards for evaluating and tracking changes in river conditions, and reviewing this report. The information presented in this report grew out of discussions involving these experts and represents the product of a collective effort; it does not reflect the opinions or viewpoints of any individual member of the technical team. The viewpoints and opinions expressed in the discussions of the group and captured in this report also do not reflect the opinions or viewpoints of the agencies, institutions, or organizations with whom the technical team members and external reviewers are associated or employed. Any errors or omissions contained herein are solely those of the Sonoran Institute.

MEMBERS OF THE LIVING RIVER TECHNICAL COMMITTEE

Plácido Dos Santos, retired water expert and member of the public

Jennifer Duan, University of Arizona

Edward Curley, Pima County Regional Wastewater Reclamation Department (retired)

Eve Halper, Bureau of Reclamation

Akitsu Kimoto, Stantec

John Kmiec, Town of Marana (formerly)

Kendall Kroesen, Tucson Audubon Society (formerly)

Michael F. Liberti, City of Tucson, Water Department

Christopher Magirl, U. S. Geological Survey

Jean E. McLain, University of Arizona

Brian Powell, Pima County Office of Sustainability and Conservation

E. Linwood Smith, Consulting Ecologist (in memoriam)

Patrice Spindler, Arizona Department of Environmental Quality

Juliet Stromberg, Arizona State University

Robert Webb, University of Arizona (retired)

Claire Zucker, University of Arizona

LIVING RIVER PROJECT TEAM

Evan Canfield, Pima County Regional Flood Control District

James DuBois and **James Brown**, Pima County Regional Wastewater Reclamation Department

Julia Fonseca and **Brian Powell**, Pima County Office of Sustainability and Conservation

Wendy Burroughs and **Yajaira Gray**, Pima County Natural Resources, Parks and Recreation

Claire A. Zugmeyer, **Luke Cole**, and **Amanda Smith**, Sonoran Institute

PRODUCTION CREDITS

Data synthesis, writing, and production: **Claire A. Zugmeyer**, **Amanda Smith**, and **James Brown**

Editing: **Audrey Spillane**

Charts: **Claire A. Zugmeyer**

Cover Design: **Terry Moody**

Design Template: **Terry Moody**

Image Credits: **cover left**: Spot-winged glider dragonfly (*Pantala hymenaea*) by Michael T. Bogan; **right**: Santa Cruz River by Charlie Aolokoy: www.alokoyphotographics.com **2**: Santa Cruz River and bicycle riders, ©Bill Hatcher/Sonoran Institute, 2019; artwork by Alana Torres, age 11, DeGrazia Elementary School — Robert Mayer. **3**: cloudy and clear water photos by Jennifer Duan; Gila topminnow by Bruce D. Taubert.



SONORAN
INSTITUTE

100 N. Stone Ave., Suite 400
Tucson, Arizona 85701

NON PROFIT
U.S. Postage
PAID
Permit #1454
Tucson, AZ 85701

SONORAN INSTITUTE has worked since our founding in 1990 to realize our vision that the Santa Cruz River, from Mexico to Marana, is a living, flowing river and the foundation of community health and prosperity.

The Sonoran Institute's mission is to connect people and communities with the natural resources that nourish and sustain them. We work at the nexus of commerce, community, and conservation to help people in the North American West build the communities they want to live in while preserving the values which brought them here. We envision a West where civil dialogue and collaboration are hallmarks of decision making, where people and wildlife live in harmony, and where clean water, air, and energy are assured.




Sonoran Institute



@sonoraninst




SonoranInstitute



Brianna Brock, age 11, DeGrazia Elementary School — Aaron Hart

Hayden Hishaw, age 11, Basis Tucson North — Carrie Vonier



Conserve Water By:
collect water from rain and feed it to the plants that don't get any water.

WE WANT YOUR THOUGHTS ON RIVER MANAGEMENT!

Sign up for information about upcoming workshops at: www.tiny.cc/scrworkshop

PIMA COUNTY

Pima County Regional Flood Control District
Pima County Regional Wastewater Reclamation Department
Pima County Office of Sustainability and Conservation
Pima County Natural Resources, Parks and Recreation

www.pima.gov

PIMA COUNTY BOARD OF SUPERVISORS

Richard Elías, Chairman, District 5
Ally Miller, District 1
Ramón Valadez, District 2
Sharon Bronson, District 3
Steve Christy, District 4

COUNTY ADMINISTRATOR

Chuck Huckelberry



CleanWatts



Energy
CERTIFIED

Printed with 100% New Wind Energy